

L1 ANSWER 7 OF 10 WPIX COPYRIGHT 2007 THE THOMSON CORP on STN
 ACCESSION NUMBER: 2000-172214 [16] WPIX
 DOC. NO. CPI: C2000-053751 [16]
 TITLE: New pyrimidinyl-substituted fused pyrazole
 derivatives,
 used for treating cardiovascular disorders such
 as
 hypertension, thromboembolic disease or ischemia
 B02; B03
 DERWENT CLASS:
 INVENTOR: ALONSO-ALIJA C; DEMBOWSKY K; FEURER A; HUETTER J;
 PERZBORN E; STAHL E; STASCH J; STRAUB A
 (FARB-C) BAYER AG; (FARB-C) BAYER HEALTHCARE AG
 PATENT ASSIGNEE:
 COUNTRY COUNT: 85

PATENT INFO ABBR.:

PATENT NO	KIND	DATE	WEEK	LA	PG	MAIN IPC
DE 19834047	A1	20000203	(200016)*	DE	35[0]	
<--						
WO 2000006568	A1	20000210	(200016)	DE		
AU 9952839	A	20000221	(200029)	EN		
EP 1102767	A1	20010530	(200131)	DE		
JP 2002521482	W	20020716	(200261)	JA	116	
US 6833364	B1	20041221	(200501)	EN		
EP 1102767	B1	20051102	(200574)	DE		
DE 59912742	G	20051208	(200581)	DE		
ES 2251213	T3	20060416	(200631)	ES		

APPLICATION DETAILS:

PATENT NO	KIND	APPLICATION	DATE
-			
DE 19834047 A1		***DE 1998-19834047	
19980729***			
AU 9952839 A		AU 1999-52839	19990716
DE 59912742 G		DE 1999-512742	19990716
EP 1102767 A1		EP 1999-938272	19990716
EP 1102767 B1		EP 1999-938272	19990716
DE 59912742 G		EP 1999-938272	19990716
WO 2000006568 A1		WO 1999-EP5073	19990716
EP 1102767 A1		WO 1999-EP5073	19990716
JP 2002521482 W		WO 1999-EP5073	19990716
US 6833364 B1		WO 1999-EP5073	19990716
EP 1102767 B1		WO 1999-EP5073	19990716
DE 59912742 G		WO 1999-EP5073	19990716
JP 2002521482 W		JP 2000-562370	19990716
US 6833364 B1		US 2001-744703	20010326
ES 2251213 T3		EP 1999-938272	19990716

FILING DETAILS:

PATENT NO	KIND	PATENT NO
DE 59912742	G Based on	EP 1102767 A

AU 9952839	A	Based on	WO 2000006568	A
EP 1102767	A1	Based on	WO 2000006568	A
JP 2002521482	W	Based on	WO 2000006568	A
US 6833364	B1	Based on	WO 2000006568	A
EP 1102767	B1	Based on	WO 2000006568	A
DE 59912742	G	Based on	WO 2000006568	A
ES 2251213	T3	Based on	EP 1102767	A

PRIORITY APPLN. INFO: DE 1998-19834047 19980729

AN 2000-172214 [16] WPIX

AB DE 19834047 A1 UPAB: 20060518

NOVELTY - 1-(Cyclic substituted methyl) 3-(cycloalkyl-substituted 2-pyrimidinyl) 4,5-fused pyrazole derivatives (I) are new.

DETAILED DESCRIPTION - Pyrazole derivatives of formula (I)

and

their isomers and salts are new.

At least one of R1, X, Y = saturated or partially

unsaturated 3-8C

cycloalkyl (optionally substituted (os) by one or more of NH2, N3,

SH,

COOH, OH, morpholino, piperidino, pyrrolidino, acyl, acylamino,

alkoxy,

alkylamino, up to 6C dialkylamino, alkylsulfonyl, alkylthio, up to

6C

alkoxycarbonyl, NO2, CN, halo, Ph, or alkyl or cycloalkyl

(optionally

substituted by NH2, SH, COOH, morpholino, piperidino, pyrrolidino,

acyl,

acylamino, alkoxy, alkylamino, up to 6C dialkylamino,

alkylsulfonyl,

alkylthio, Ph, alkylsulfonylamino, up to 6C alkoxycarbonyl, NO2,

CN or

halo));

any other R1, X, Y = H or a very wide range of specific substituents;

R2 + R3 = group completing a phenyl ring or a 6-membered

saturated

or aromatic heterocycle (containing 1-3 of N, O and S), optionally

substituted by a wide range of specific groups;

A = 5- or 6-membered saturated or aromatic heterocycle

(containing

1-3 of N, O and S), optionally substituted by a wide range of

specific

groups;

unless specified otherwise alkyl, acyl and cycloalkyl

moieties have

up to 6C. Full definitions are given in the DEFINITIONS.

An INDEPENDENT CLAIM is included for the preparation of

(I).

ACTIVITY - Cardiovascular; vascular relaxant;

antithrombotic;

hypotensive; coronary dilator; antiangina; antiarrhythmic;

antiischemic;

urogenital; neuroprotective; anxiolytic; antidepressant;

analgesic.

3-(4-Amino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo

(3,4-b) pyridine (Ia) at 1 mg/kg p.o. gave a maximum blood pressure decrease of 23 mm Hg after 20 minutes in narcotized rats.
MECHANISM OF ACTION - Soluble guanyl cyclase stimulant; intracellular cyclic guanosine monophosphate (cGMP) level increasing agent. (I) also potentiates the activity of other agents which increase cGMP levels, e.g. endothelium derived relaxing factor (EDRF), nitric oxide donors, protoporphyrin IX, arachidonic acid or phenylhydrazine derivatives.
USE - (I) cause vascular relaxation, inhibit thrombocyte aggregation, reduce blood pressure and increase coronary blood flow. They are used for treating cardiovascular disorders (claimed), e.g. hypertension, cardiac insufficiency, angina pectoris, peripheral or cardiac vascular disease, arrhythmia, thromboembolic disease or ischemia (claimed) (e.g. myocardial infarction, cerebral stroke, transitory ischemic attacks, peripheral blood flow disorders or restenosis), arteriosclerosis or diseases of the urogenital system (e.g. prostate hypertrophy, erectile dysfunction, female sexual dysfunction or incontinence). (I) are also useful for treating central nervous system disorders, especially for alleviating cognitive deficiency, improving learning and memory performance or treating Alzheimer's disease but also for treating anxiety, stress, depression, CNS-related sexual dysfunction, sleep disorders or food, condiment and sweetener uptake disorders. (I) are further useful for regulating cerebral blood flow, treating migraine or pain and treating or preventing the sequelae of cerebral infarction (e.g. stroke), cerebral ischemia or cranial-cerebral trauma.

Member(0002)

ABEQ WO 2000006568 Al UPAB 20060518

NOVELTY - 1-(Cyclic substituted methyl) 3-(cycloalkyl-substituted 2-pyrimidinyl) 4,5-fused pyrazole derivatives (I) are new.

DETAILED DESCRIPTION - Pyrazole derivatives of formula (I) and their isomers and salts are new.

At least one of R1, X, Y = saturated or partially unsaturated 3-8C

cycloalkyl (optionally substituted (os) by one or more of NH2, N3, SH,

COOH, OH, morpholino, piperidino, pyrrolidino, acyl, acylamino, alkoxy,

alkylamino, up to 6C dialkylamino, alkylsulfonyl, alkylthio, up to 6C

alkoxycarbonyl, NO₂, CN, halo, Ph, or alkyl or cycloalkyl
(optionally substituted by NH₂, SH, COOH, morpholino, piperidino, pyrrolidino,
acyl,
acylamino, alkoxy, alkylamino, up to 6C dialkylamino,
alkylsulfonyl,
alkylthio, Ph, alkylsulfonylamino, up to 6C alkoxycarbonyl, NO₂,
CN or

halo));
any other R₁, X, Y = H or a very wide range of specific
substituents;

R₂ + R₃ = group completing a phenyl ring or a 6-membered
saturated
or aromatic heterocycle (containing 1-3 of N, O and S), optionally
substituted by a wide range of specific groups;

A = 5- or 6-membered saturated or aromatic heterocycle
(containing
1-3 of N, O and S), optionally substituted by a wide range of
specific
groups;

unless specified otherwise alkyl, acyl and cycloalkyl
moieties have

up to 6C. Full definitions are given in the DEFINITIONS.

An INDEPENDENT CLAIM is included for the preparation of
(I).

ACTIVITY - Cardiovascular; vascular relaxant;
antithrombotic;
hypotensive; coronary dilator; antiangina; antiarrhythmic;
antiischemic;
urogenital; neuroprotective; anxiolytic; antidepressant;
analgesic.

3-(4-Amino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-
pyrazolo

(3,4-b) pyridine (Ia) at 1 mg/kg p.o. gave a maximum blood
pressure

decrease of 23 mm Hg after 20 minutes in narcotized rats.

MECHANISM OF ACTION - Soluble guanyl cyclase stimulant;
intracellular cyclic guanosine monophosphate (cGMP) level
increasing

agent. (I) also potentiates the activity of other agents which
increase

cGMP levels, e.g. endothelium derived relaxing factor (EDRF),
nitrogen

monoxide donors, protoporphyrin IX, arachidonic acid or
phenylhydrazine
derivatives.

USE - (I) cause vascular relaxation, inhibit thrombocyte
aggregation, reduce blood pressure and increase coronary blood
flow. They

are used for treating cardiovascular disorders (claimed), e.g.
hypertension, cardiac insufficiency, angina pectoris, peripheral
or

cardiac vascular disease, arrhythmia, thromboembolic disease or
ischemia

(claimed) (e.g. myocardial infarction, cerebral stroke, transitory
ischemic attacks, peripheral blood flow disorders or restenosis),

arteriosclerosis or diseases of the urogenital system (e.g. prostate hypertrophy, erectile dysfunction, female sexual dysfunction or incontinence). (I) are also useful for treating central nervous system disorders, especially for alleviating cognitive deficiency, improving learning and memory performance or treating Alzheimer's disease but also for treating anxiety, stress, depression, CNS-related sexual dysfunction, sleep disorders or food, condiment and sweetener uptake disorders. (I) are further useful for regulating cerebral blood flow, treating migraine or pain and treating or preventing the sequelae of cerebral infarction (e.g. stroke), cerebral ischemia or cranial-cerebral trauma.

Member(0004)

ABEQ EP 1102767 A1 UPAB 20060518

NOVELTY - 1-(Cyclic substituted methyl) 3-(cycloalkyl-substituted 2-pyrimidinyl) 4,5-fused pyrazole derivatives (I) are new.

DETAILED DESCRIPTION - Pyrazole derivatives of formula (I)

and

their isomers and salts are new.

At least one of R1, X, Y = saturated or partially

unsaturated 3-8C

cycloalkyl (optionally substituted (os) by one or more of NH2, N3,

SH,

COOH, OH, morpholino, piperidino, pyrrolidino, acyl, acylamino,

alkoxy,

alkylamino, up to 6C dialkylamino, alkylsulfonyl, alkylthio, up to

6C

alkoxycarbonyl, NO2, CN, halo, Ph, or alkyl or cycloalkyl

(optionally

substituted by NH2, SH, COOH, morpholino, piperidino, pyrrolidino,

acyl,

acylamino, alkoxy, alkylamino, up to 6C dialkylamino,

alkylsulfonyl,

alkylthio, Ph, alkylsulfonylamino, up to 6C alkoxycarbonyl, NO2,

CN or

halo));

any other R1, X, Y = H or a very wide range of specific substituents;

R2 + R3 = group completing a phenyl ring or a 6-membered

saturated

or aromatic heterocycle (containing 1-3 of N, O and S), optionally substituted by a wide range of specific groups;

A = 5- or 6-membered saturated or aromatic heterocycle

(containing

1-3 of N, O and S), optionally substituted by a wide range of specific groups;

unless specified otherwise alkyl, acyl and cycloalkyl moieties have

up to 6C. Full definitions are given in the DEFINITIONS.

An INDEPENDENT CLAIM is included for the preparation of (I).

ACTIVITY - Cardiovascular; vascular relaxant; antithrombotic; hypotensive; coronary dilator; antiangina; antiarrhythmic; antiischemic; urogenital; neuroprotective; anxiolytic; antidepressant; analgesic.

3-(4-Amino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo

(3,4-b) pyridine (Ia) at 1 mg/kg p.o. gave a maximum blood pressure

decrease of 23 mm Hg after 20 minutes in narcotized rats.

MECHANISM OF ACTION - Soluble guanyl cyclase stimulant; intracellular cyclic guanosine monophosphate (cGMP) level increasing

agent. (I) also potentiates the activity of other agents which increase

cGMP levels, e.g. endothelium derived relaxing factor (EDRF), nitrogen

monoxide donors, protoporphyrin IX, arachidonic acid or phenylhydrazine derivatives.

USE - (I) cause vascular relaxation, inhibit thrombocyte aggregation, reduce blood pressure and increase coronary blood flow. They

are used for treating cardiovascular disorders (claimed), e.g. hypertension, cardiac insufficiency, angina pectoris, peripheral or

cardiac vascular disease, arrhythmia, thromboembolic disease or ischemia

(claimed) (e.g. myocardial infarction, cerebral stroke, transitory ischemic attacks, peripheral blood flow disorders or restenosis), arteriosclerosis or diseases of the urogenital system (e.g.

prostate hypertrophy, erectile dysfunction, female sexual dysfunction or incontinence). (I) are also useful for treating central nervous

system disorders, especially for alleviating cognitive deficiency, improving

learning and memory performance or treating Alzheimer's disease but also

for treating anxiety, stress, depression, CNS-related sexual dysfunction,

sleep disorders or food, condiment and sweetener uptake disorders. (I) are

further useful for regulating cerebral blood flow, treating migraine or

pain and treating or preventing the sequelae of cerebral infarction (e.g.

stroke), cerebral ischemia or cranial-cerebral trauma.